



Atty Dkt 2300-0054.08
Client Dkt 0054.009
PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In Re Application of:

HELDIN et al.

Serial No.: 08/453,350

Group Art Unit: 1812

Filing Date: May 30, 1995

Examiner: K. Brown

Title: RECOMBINANT PDGF A-CHAIN
HOMODIMERS AND METHODS OF
USE

RECEIVED

Declaration of Christer Betsholtz, Ph.D. MAR 09 1998

Assistant Commissioner for Patents
Washington, D.C. 20231

MATRIX CUSTOMER
SERVICE CENTER

Sir:

I, Christer Betsholtz, declare as follows:

1. I am a coinventor of the above-identified application. I hold a Ph.D. in Experimental Pathology from the University of Uppsala, Sweden. I am employed by the Department of Medical Biochemistry at the University of Gothenburg, Gothenburg Sweden, as a Professor of Medical Biochemistry and have held this position since 1993. Prior to that time, I was a Senior Research Fellow at the Cancer Research Fund, Sweden and held Post doctoral positions at the Cancer Research Fund and the Medical Research Council, Sweden. A copy of my curriculum vitae is attached hereto as Exhibit A.

2. I am extremely familiar with PDGF and have actively been studying this molecule, including the molecular characterization and mechanisms of action thereof, for over 15

Atty Dkt No. 2300-0054.08
USSN: 08/453,350
PATENT

years. I have coauthored numerous publications regarding PDGF. A true and correct copy of my curriculum vitae is attached hereto as Exhibit A.

3. I have reviewed the Office Action dated July 8, 1997 ("the Action"), as well as Heldin et al., *Nature* (1986) 319:511-514 ("Heldin"), cited in the Action, on which I am a coauthor. In the Heldin paper, we describe the isolation of what was termed therein osteosarcoma-derived growth factor ("ODGF"). We now know this molecule to be the same as a naturally occurring PDGF A-chain homodimer. However, at the time of the publication, we did not realize that this molecule was the same as a naturally occurring A-chain homodimeric form of PDGF because we did not know that such a form of naturally occurring PDGF existed (see, page 513, second column, of Heldin).

4. The ODGF described in Heldin was isolated directly from a human osteosarcoma cell line using sequential chromatography of conditioned medium from U-2 osteosarcoma cells. This cell line was established from a human patient suffering from cancer. Although this cell line was propagated in culture, the cell line may well have contained pathogenic viruses from the patient. Moreover, since the cell line was of human origin, it could easily become infected with human pathogenic viruses during propagation. Therefore, I believe the Action's statement on page 4 that the ODGF "would be highly unlikely to be contaminated with virus" to be in error. The methods used to purify ODGF described in Heldin would not guarantee the elimination of human viruses. This method would not be appropriate for producing an ODGF compound to be used in pharmaceutical compositions since viral contaminants may be present. Accordingly, there would be a high

Atty Dkt No. 2300-0054.08
USSN: 08/453,350
PATENT

risk associated with using ODGF purified as described in Heldin, in compositions for treating patients.

5. Additionally, the methods described in Heldin would not produce ODGF preparations completely free of other human proteins. In particular, ODGF was isolated using a Sephacryl S-200 column, followed by a BioGel P-150 column, and then an HPLC RP8 column. The product of each of these methods would inherently include at least small amounts of human proteins other than human ODGF since the ODGF was isolated from human osteosarcoma cells. The methods for purifying proteins from human sources described above, cannot result in a protein product free of contaminating human proteins. The Action notes that amino acid sequence analysis and silver staining are highly sensitive methods for determining whether a protein sample is homogeneous. Despite these observations, it is a virtual certainty that trace amounts of human proteins were present in the ODGF preparations that were not detected by these methods.

6. Recombinant methods of producing human PDGF A-chain, such as those described in the subject application, on the other hand, result in a preparation free of other human proteins and devoid of contaminating human viruses. This is because the only human structural gene present in the recombinant plasmids is the gene encoding human PDGF. It would not be possible to produce preparations having such purity without the gene encoding PDGF. Heldin does not describe the gene or recombinant methods for producing PDGF A-chain.

Atty Dkt No. 2300-0054.08
USSN: 08/453,350
PATENT

7. I declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

98-02-11
Date

Christer Betsholtz
Christer Betsholtz, Ph.D.

EXHIBIT A

CURRICULUM VITAE

CHRISTER BETSHOLTZ

Born: July 11, 1959 in Stockholm, Sweden.

Home address: Haraldssons väg 11
421 67 Västra Frölunda

Home telephone: +46 31 286459

Work address: Department of Medical Biochemistry
University of Gothenburg
Medicinargatan 9A
SE 405 30 Gothenburg
Sweden

Telephone: +46 31 7733460 (office) 7733463 (lab)

Telefax: +46 31 416108

E-mail: Christer.Betsholtz@medkem.gu.se

Citizenship: Swedish

Family Married to Ingrid Holmberg-Betsholtz, two children born 1988 and 1990

Degrees and appointments (in chronological order):

1980	Bachelor of Medicine, University of Uppsala
1986	PhD
1986-88	Post doctoral position (The Cancer Research Fund, Sweden)
1989-90	Post doctoral position (The Medical Research Council, Sweden)
1991-93	Senior Research Fellow (The Cancer Research Fund, Sweden)
1993-	Professor of Medical Biochemistry, Gothenburg University, Sweden

Scientific awards

1989: The Oscar Prize, Uppsala University
1995: Fernström's Prize for young investigators, Gothenburg University
1997: Göran Gustavsson's Award in Molecular Biology

Scientific projects

Genetic analysis of platelet-derived growth factor functions.
Islet amyloid polypeptide (amylin) molecular genetics.
Glial fibrillary acidic protein, intermediate filaments and astrocyte functions.

Invited lectures at international symposia (from 1994-1997, selected):

Keystone symposium "Biology of the vascular wall/endothelial cells" Keystone, CO 1994
Second Nordic workshop on transgenic mice, Turku, Finland 1994
Reaction to injury revisited, Univ of Washington Symposium, Seattle, WA 1994
Tsumagoi Conference on cardiovascular biology, Kyoto, Japan, 1995
International Vascular Biology Meeting, Seattle, WA 1996
ISN Forefronts in Nephrology, Snowbird, Utah, 1996
Keystone symposium "Inflammation, Growth regulatory molecules and Atherosclerosis", Keystone CO 1997
Cold Spring Harbor symposium on Signal transduction in endothelial cells 1997

1998: Gordon Conference on vascular biology, July 1998
Juselius Symposium on angiogenesis, Helsinki June 1998
FEBS Silver Jubilee meeting, Copenhagen July 1998

Morphogenesis of the endothelium, Schloss Ringberg Germany, September 1998
7th International workshop Developmental Biology, Stockholm September 1998
5th Franz-Volhard Symposium, Berlin Germany, September 1998

Ad-hoc reviewer for:

Atherosclerosis
Cancer Research
Development
Diabetologia
European Journal of Biochemistry
Experimental Cell Research
FEBS Letters
Growth Factors
International Journal of Cancer
Journal of Biological Chemistry
Journal of Cell Biology
Nature
Nucleic Acids Research
The Cancer Journal

Scientific publications (selected)

Betsholtz, C., Heldin, C.-H., Nistér, M., Ek, B., Wasteson, Å. and Westermark, B. Synthesis of a PDGF-like growth factor in human glioma and sarcoma cells suggests the expression of the cellular homologue to the transforming protein of simian sarcoma virus. *Biochem Biophys Res Commun* **117**, 176-182 (1983)

Betsholtz, C. and Westermark, B. Growth factor-induced proliferation of human fibroblasts in serum-free culture depends on cell density and extracellular calcium concentration. *J Cell Physiol* **118**, 203-210 (1984)

Betsholtz, C., Westermark, B., Ek, B. and Heldin, C.-H. Coexpression of a PDGF-like growth factor and PDGF receptors in a human osteosarcoma cell line: Implications for autocrine receptor activation. *Cell* **39**, 447-457 (1984)

Johnsson, A., Betsholtz, C., von der Helm, K., Heldin, C.-H. and Westermark, B. Platelet-derived growth factor agonist activity of a secreted form of the v-sis oncogene product. *Proc Natl Acad Sci USA* **82**, 1721-1725 (1985)

Goustin, A.S., Betsholtz, C., Pfeifer-Ohlsson, S., Persson, H., Rydner, J., Bywater, M., Holmgren, G., Heldin, C.-H., Westermark, B. and Ohlsson, R. Coexpression of the *sis* and *myc* proto-oncogenes in developing human placenta suggests autocrine control of trophoblast growth. *Cell* **41**, 301-312 (1985)

Pfeifer-Ohlsson, S., Rydner, J., Goustin, A.S., Larsson, E., Betsholtz, C. and Ohlsson, R. Cell-type-specific pattern of *myc* proto-oncogene expression in developing human embryos. *Proc Natl Acad Sci USA* **82**, 5050-5054 (1985)

Betsholtz, C., Bywater, M., Westermark, B., Bürk, R. and Heldin, C.-H. Expression of the *c-sis* gene and secretion of a platelet-derived growth factor-like molecule by simian virus 40-transformed BHK cells. *Biochem Biophys Res Commun* **130**, 753-760 (1985)

Johnsson, A., Betsholtz, C., Heldin, C.-H. and Westermark, B. Antibodies against platelet-derived growth factor inhibit acute transformation by simian sarcoma virus. *Nature* **317**, 438-440 (1985)

Knott, T.J., Rall, S.C., Innerarity, T.L., Jacobson, S.F., Urdea, M.S., Levy-Wilson, B., Powell, L., Pease, R., Eddy, R., Nakai, H., Byers, M., Priestley, L., Robertson, E., Rall, L.B., Betsholtz, C., Shows, T.B., Mahley, R.W. and Scott, J. Human apolipoprotein B: Structure of carboxyterminal domains, sites of gene expression and chromosomal localization. *Science* **220**, 37-43 (1985)

Heldin, C.-H., Johnsson, A., Wennergren, S., Wernstedt, C., Betsholtz, C. and Westermark, B. A human osteosarcoma cell line secretes a growth factor structurally related to a homodimer of PDGF A chains. *Nature* **319**, 511-514 (1986)

Betsholtz, C., Johnsson, A., Heldin, C.-H., Westermark, B., Lind, P., Urdea, M.S., Shows, T.B., Philpott, K., Mellor, A., Knott, T.J. and Scott, J. Complementary DNA sequence and chromosomal

localisation of human platelet-derived growth factor A-chain and its expression in tumour cell lines. *Nature* **320**,695-699 (1986)

Johnsson, A., Betsholtz, C., Heldin, C.-H. and Westermark, B. The phenotypic characteristics of simian sarcoma virus-transformed human fibroblasts suggest that the *v-sis* gene product acts solely as a PDGF receptor agonist in cell transformation. *EMBO J* **5**,1535-1541 (1986)

Paulsson, Y., Bywater, M., Pfeifer-Ohlsson, S., Ohlsson, R., Heldin, C.-H., Westermark, B. and Betsholtz, C. Growth factors induce early pre-replicative changes in senescent human fibroblasts. *EMBO J* **5**,2157-2162 (1986)

Betsholtz, C., Johnsson, A., Heldin, C.-H. and Westermark, B. Efficient reversion of simian sarcoma virus-transformation and inhibition of growth factor-induced mitogenesis by suramin. *Proc Natl Acad Sci USA* **83**,6440-6444 (1986)

Sejersen T., Betsholtz, C., Sjölund, M., Heldin, C.-H., Westermark, B. and Thyberg, J. Rat skeletal myoblasts and arterial smooth muscle cells express the gene for the A-chain but not the B-chain (c-sis) of platelet-derived growth factor (PDGF) and produce a PDGF-like protein. *Proc Natl Acad Sci USA* **83**,6844-6848 (1986)

Westermark, B., Johnsson, A., Paulsson, Y., Betsholtz, C., Heldin, C.-H., Herlyn, M., Rodeck, U. and Koprowski, H. Human melanoma cell lines of primary and metastatic origin express the genes encoding the chains of platelet-derived growth factor (PDGF) and produce a PDGF-like growth factor. *Proc Natl Acad Sci USA* **83**, 7197-7200 (1986)

Betsholtz, C., Bergh, J., Bywater, M., Pettersson, M., Johnsson, A., Heldin, C.-H., Ohlsson, R., Knott, T.J., Scott, J., Bell, G.I. and Westermark, B. Expression of multiple growth factors in a human lung cancer cell line. *Int J Cancer* **39**,502-507 (1987)

Collins, T., Pober, J.S., Gimbrone Jr, M.A., Hammacher, A., Betsholtz, C., Westermark, B. and Heldin, C.-H. Cultured human endothelial cells express platelet-derived growth factor A-chain. *Am J Pathol* **127**,7-12 (1987)

Alitalo, R., Andersson, L.C., Betsholtz, C., Nilsson, K., Westermark, B., Heldin, C.-H. and Alitalo, K. Induction of platelet-derived growth factor gene expression during megakaryoblastic and monocytic differentiation of human leukemia cell lines. *EMBO J* **6**,1213-1218 (1987)

Peres, R., Betsholtz, C., Westermark, B. and Heldin, C.-H. Frequent expression of growth factors for mesenchymal cells in human mammary carcinoma cell lines. *Cancer Res* **47**,3425-3429 (1987)

Nistér, M., Wedell, B., Betsholtz, C., Bywater, M., Pettersson, M., Westermark, B. and Mark, G. Evidence for progression changes in the human malignant glioma line U-343 MG: Analysis of karyotype and expression of genes encoding the subunit chains of platelet-derived growth factor. *Cancer Res.* **47**,4953-4961 (1987)

Rorsman, F., Bywater, M., Knott, T.J., Scott, J. and Betsholtz, C. Structural characterization of the platelet-derived growth factor A-chain cDNA and gene: Alternative exon usage predicts two different precursor proteins. *Mol Cell Biol* **8**,571-577 (1988)

Söderdahl, G., Betsholtz, C., Johansson, A., Nilsson, K. and Bergh, J. Differential expression of platelet-derived growth factor and transforming growth factor genes in small- and non-small cell human lung cancer cell lines. *Int J Cancer* **41**,636-641 (1988)

Bywater, M., Rorsman, F., Bongcam-Rudloff, E., Mark, G., Heldin, C.-H., Westermark, B. and Betsholtz, C. Expression of recombinant platelet-derived growth factor A- and B-chain homodimers in Rat-1 cells and human fibroblasts reveals differences in protein processing and autocrine effects. *Mol Cell Biol* **8**,2753-2762. (1988)

Nistér, M., Libermann, T.A., Betsholtz, C., Pettersson, M., Claesson-Welsh, L., Heldin, C.-H., Schlessinger, J. and Westermark, B. Expression of messenger RNAs for platelet-derived growth factor and transforming growth factor- α and their receptors in human glioma cell lines. *Cancer Res* **48**,3910-3918 (1988)

Claesson-Welsh, L., Eriksson, A., Morén, A., Severinsson, L., Ek, B., Östman, A., Betsholtz, C. and Heldin, C.-H. cDNA cloning and expression of a human platelet-derived growth factor (PDGF) receptor specific for B-chain-containing PDGF molecules. *Mol Cell Biol* **8**,3476-3486 (1988)

- Beckmann, M.P., Betsholtz, C., Heldin, C.-H., Westermark, B., Di Marco E., Di Fiore, P.P. Robbins, K.C., and Aaronson, S.A. Comparison of biological properties and transforming potential of human PDGF-A and PDGF-B chains. *Science* **241**, 1346-1349 (1988)
- Stenman, G., Rorsman, F. and Betsholtz, C. Sublocalization of the PDGF A-chain gene to human chromosome 7, band q11.23 by *in situ* hybridization. *Exp Cell Res* **178**, 180-184 (1988)
- Östman, A., Rull, L., Hammocher, A., Wormsted, M.A., Coit, D., Valenzuela, P., Betsholtz, C., Westermark, B., and Heldin, C.-H. Synthesis and assembly of a functionally active recombinant platelet-derived growth factor AB heterodimer. *J Biol Chem* **263**, 16202-16208 (1988)
- Hermansson, M., Nistér, M., Betsholtz, C., Heldin, C.-H., Westermark, B. and Funa, K. Endothelial cell hyperplasia in human glioblastoma: coexpression of mRNA for platelet-derived growth factor β chain and PDGF receptor suggests autocrine growth stimulation. *Proc Natl Acad. Sci USA* **85**, 7748-7752 (1988)
- Ashcroft, F.M., Ashcroft, S.J.H., Berggren, P.-O., Betsholtz, C., Rorsman, P., Trube, G. and Welsh, M. Expression of K-channels in *Xenopus laevis* oocytes injected with poly(A⁺) RNA from the insulin-secreting β -cell line HIT T15. *FEBS Lett* **239**, 185-189 (1988)
- Betsholtz, C., Johnson, K.H. and Westermark, P. "Amylin" hormone (Scientific Correspondence) *Nature* **338**, 221 (1989)
- Matoskova, B., Rorsman, F., Svensson, V. and Betsholtz, C. Alternative splicing of the platelet-derived growth factor A chain transcript: Occurrence in normal as well as transformed cells and in several mammalian species. *Mol Cell Biol* **9**, 3148-3150 (1989)
- Betsholtz, C., Christmansson, L., Engström, U., Rorsman, F., Svensson, V., Johnson, K.H. and Westermark, P. Sequence divergence in a specific region of islet amyloid polypeptide (IAPP) explains differences in islet amyloid formation between species. *FEBS Lett* **251**, 261-264 (1989)
- Östman, A., Bäckström, G., Fong, N., Betsholtz, C., Wernstedt, C., Hellman, U., Westermark, B., Valenzuela, P. and Heldin, C.-H. Expression of three recombinant homodimeric isoforms of PDGF in *Saccharomyces cerevisiae*. Evidence for differences in receptor binding and functional activities. *Growth factors* **1**, 271-281 (1989)
- Betsholtz, C., Svensson, V., Rorsman, F., Engström, U., Westermark, G.T., Wilander, E., Johnson, K.H. and Westermark, P. Islet amyloid polypeptide: cDNA cloning and identification of an amyloidogenic region associated with the species-specific occurrence of age-related diabetes mellitus. *Exp Cell Res* **183**, 484-493 (1989)
- Westermark, P., Engström, U., Westermark, G.T., Johnson, K.H., Permerth, J. and Betsholtz, C. Islet amyloid polypeptide (IAPP) and pro-IAPP immunoreactivity in human islets of Langerhans. *Diabetes Res Clin Pract* **7**, 219-226 (1989)
- Johnson, K.H., O'Brien, T.D., Betsholtz, C. and Westermark, P. Islet amyloid, islet amyloid polypeptide and diabetes mellitus. *New Engl J Med* **321**, 513-518 (1989)
- Betsholtz, C., Christmansson, L., Engström, U., Rorsman, F., Jordan, K., O'Brien, T., Murtaugh, M., Johnson, K.H. and Westermark, P. The structure of cat islet amyloid polypeptide (IAPP) and identification of amino acid residues of potential significance for islet amyloid formation. *Diabetes* **39**, 118-122 (1990)
- Johnson, K.H., O'Brien, T.D., Jordan, K., Betsholtz, C. and Westermark, P. The putative hormone islet amyloid polypeptide (IAPP) induces impaired glucose tolerance in cats. *Biochem Biophys Res Commun* **167**, 507-513 (1990)
- Betsholtz, C., Baumann, A., Ashcroft, F.M., Ashcroft, S.J.H., Berggren, P.-O., Pongs, O., Rorsman, P., Sandblom, J., and Welsh, M. Expression of voltage-gated channels in insulin-producing cells: Analysis by polymerase chain reaction. *FEBS Lett* **263**, 121-126 (1990)
- Betsholtz, C., Rorsman, F., Westermark, B., Östman, A. and Heldin, C.-H. Analogous alternative splicing (Scientific correspondence) *Nature* **344**, 299 (1990)
- Westermark, P., Engström, U., Westermark, G.T., Johnson, K.H. and Betsholtz, C. Islet amyloid polypeptide: Pinpointing amino acid residues linked to amyloid fibril formation. *Proc Natl Acad Sci USA* **87**, 5036-5040 (1990)

Christmanson, L., Rorsman, F., Stenman, G., Westermark, P. and Betsholtz, C. The human islet amyloid polypeptide gene: Organization, chromosomal localization and identification of a promoter region. *FEBS Lett.* **267**, 160-166 (1990)

Jordan, K., Murtaugh, M.P., O'Brien, T.D., Westermark, P., Betsholtz, C. and Johnson, K.H. cDNA predicted sequence of canine IAPP provides important clues regarding pathogenesis and amyloidogenesis in type 2 diabetes. *Biochem Biophys Res Commun* **169**, 502-508. (1990)

Welsh, M., Claesson-Welsh, L., Hallberg, A., Welsh, N., Betsholtz, C., Arkhammar, P., Nilsson, T., Heldin, C.-H. and Berggren, P.-O. Coexpression of the platelet-derived growth factor (PDGF) B chain and the PDGF β receptor in isolated pancreatic islet cells stimulates DNA synthesis. *Proc Natl Acad Sci USA* **87**, 5807-5811 (1990)

Leveen, P., Claesson, Welsh, L., Heldin, C.-H., Westermark, B. and Betsholtz, C. Expression of the genes for PDGFs and its receptors in human sarcoma cell lines. *Int J Cancer* **46**, 1066-1070 (1990).

Madsen, O., Nielsen, J.H., Michelsen, B., Westermark, P., Betsholtz, C., Nishi, M. and Steiner, D.F. Islet amyloid polypeptide (IAPP/Amylin) and insulin expression are controlled differently in primary and transformed islet cells. *Mol Endo* **5**, 143-148 (1991)

Östman, A., Andersson, M., Betsholtz, C., Westermark, B. and Heldin, C.-H. Identification of a cell retention signal in the B-chain of PDGF and in the long splice version of the A-chain. *Cell Regul.* **2**, 503-512 (1991)

Bongcam-Rudloff, E., Nistér, M., Stenman, G., Huebner, K., Croce, C.M., Betsholtz, C. and Westermark, B. Human glial fibrillary acidic protein: Complementary DNA cloning, chromosomal localization and messenger RNA expression in human glioma cell lines of various phenotypes. *Cancer Res* **51**, 1553-1560 (1991).

Christmanson, L., Betsholtz, C., Johansson, B. and Westermark, P. The transthyretin cDNA sequence is normal in transthyretin-derived senile systemic amyloidosis. *FEBS Lett* **281**, 177-180 (1991).

Zierath, J.R., Galuska, D., Engström, Å., Johnson, K.H., Betsholtz, C., Westermark, P. and Wallberg-Henriksson, H. Synthetic human islet amyloid polypeptide inhibits insulin and phorbol ester-stimulated glucose transport in in vitro incubated human muscle strips. *Diabetologia* **35**, 26-31 (1992).

Rorsman, F. and Betsholtz, C. Characterization of the mouse PDGF A chain gene. Evolutionary conservation of gene structure, nucleotide sequence and alternative splicing. *Growth Factors* **6**, 303-313 (1992)

Westermark, P., Johnson, K.H. and Betsholtz, C. Islet amyloid polypeptide - a novel controversy in diabetes research. *Diabetologia* **35**, 297-303 (1992)

Stenman, G., Rorsman, F., Huebner, K. and Betsholtz, C. The human platelet-derived growth factor A chain gene (PDGFA) maps to chromosome 7p22. *Cytogenet. Cell. Genet.* **60**, 206-207 (1992)

Rorsman, F., Leveen, P. and Betsholtz, C. Platelet-derived growth factor (PDGF) A-chain heterogeneity generated by the use of alternative promoters and alternative polyadenylation sites. *Growth Factors* **7**, 241-251 (1992).

Johnson, K.H., O'Brien, T.D., Betsholtz, C. and Westermark, P. Islet amyloid polypeptide (IAPP): Mechanisms of amyloidogenesis in pancreatic islets and potential roles in diabetes mellitus. *Lab. Invest* **66**, 522-535 (1992)

Leveen, P., Betsholtz, C. and Westermark, B. Evidence for negative transacting mechanisms controlling expression of platelet-derived growth factor A-chain and B-chain mRNA in somatic cell hybrids. *Experimental Cell Res.* **207**, 283-289 (1993)

Örtoft, E., Pählman, S., Andersson, G., Parrow, V., Betsholtz, C. and Hammerling, U. Human GAP-43 gene expression: Multiple start sites for initiation of transcription in differentiating neuroblastoma cells. *Mol. Cell. Neurosci.* in press (1993)

Christmanson, L., Betsholtz, C., Leckström, A., Engström, U., Cortie, C., Johnson, K.H. and Westermark, P. Islet amyloid polypeptide in the rabbit and European hare: Relationship with islet amyloidogenesis. *Diabetologia* **36**, 183-188 (1993)

- Van der Meeren, A., Seddon, M.B., Betsholtz, C., Lechner, J.F. and Gerwin, B. Tumorigenic conversion of human mesothelial cells is a consequence of platelet-derived growth factor-A chain overexpression. *Am. J. Respir. Cell Mol. Biol.* **8**, 214-221 (1993).
- Westermarck G.T., Christmansson, L., Terenghi, G., Permerth, J., Betsholtz, C., Larsson, J., Polak, J.M. and Westermarck, P. Islet amyloid polypeptide: demonstration of mRNA in human pancreatic islets by in situ hybridization and evidence of altered secretion in conjunction with islet amyloid. *Diabetologia*, **36**, 323-328 (1993).
- Velloso, L., Kümpe, O., Hallberg, A., Christmansson, L., Betsholtz, C., Karlsson, F.A. Determination of autoantibodies against glutamate decarboxylase in insulin-dependent diabetes mellitus using a radioligand produced by eukaryotic expression. *J. Clin. Invest.* **91**, 2084-2090 (1993).
- Betsholtz, C., Christmansson, L., Gebre-Medhin, S. and Westermarck, P. Islet amyloid polypeptide - hen or egg in type 2 diabetes pathogenesis? *Acta Oncologica* **32**, 149-154 (1993)
- Christmansson, L., Westermarck, P. and Betsholtz, C. Islet amyloid polypeptide (IAPP/amylin) stimulates cyclic AMP accumulation via the calcitonin receptor. *Biochem. Biophys. Res. Commun.* **205**, 1226-1235 (1994)
- Levéen, P., Pekny, P., Gebre-Medhin, S., Swolin, B., Larsson, E. and Betsholtz, C. Mice deficient for PDGF B show renal, cardiovascular and hematological abnormalities. *Genes Dev* **8**, 1875-1887 (1994)
- Kulkarni, P.S., Sundqvist, K., Betsholtz, C., Höglund, P., Wiman, K., Zhivotovsky, B., Bertolero, F., Liu, Y. and Grafström, R. Characterization of human buccal epithelial cells transfected with the simian virus 40 T-antigen gene. *Carcinogenesis* **16**, 2515-2521 (1995)
- Pekny, M., Levéen, P., Pekna, M., Eliasson, C., Westermarck, B. and Betsholtz, C. Mice lacking glial fibrillary acidic protein (GFAP) develop and reproduce normally. *EMBO J* **14**, 1590-1598 (1995)
- Pekny, M., Östman, A., Pekna, M., Törnelli, J., Feinstein, R., Heldin, C.-H., Westermarck, B. and Betsholtz, C. Luteal failure in transgenic mice carrying a PDGF dominant-negative mutant/GH hybrid transgene. *Transgenic* **1**, 515-523 (1995)
- Betsholtz, C. Role of platelet-derived growth factors in mouse development. *Int. J. Dev. Biol.* **39**, 817-825 (1995).
- Boström, H., Willetts, K., Pekny, M., Levéen, P., Hedstrand, H., Pekna, M., Schalling, M., Nilsson, M., Gebre-Medhin, S., Kurland, S., Törnelli, J., Heath, J.K. and Betsholtz, C. PDGF-A signaling is a critical event in lung alveolar myofibroblast ontogeny and alveogenesis. *Cell* **85**, 863-873 (1996)
- Lagberg Arkblad, E., Betsholtz, C., and Rydström, J. The cDNA sequence of proton-pumping nicotinamide nucleotide transhydrogenase from man and mouse. *Biochim. Biophys. Acta* **1273**, 203-205 (1996)
- Tatzelt, J., Maeda, N., Pekny, M., Yang, S.-L., Betsholtz, C., Eliasson, C., Cayetano, J., Camarino, A.P., DeArmond, S.J. and Prusiner, S. Scrapie in mice deficient for either apolipoprotein E or glial fibrillary acidic protein. *Neurology*, **47**, 449-453 (1996)
- Pekny, M., Eliasson, C., Chien, C.-L., Kindblom, L.G., Liem, R., Hamberger, A. and Betsholtz, C. GFAP-deficient astrocytes in vitro are capable of stellation when cocultured with neurons and exhibit reduced amount of intermediate filaments and increased cell saturation density. *Exp. Cell Res.* in press (1997)
- Betsholtz, C. and Raines, E.W. Platelet-derived growth factor - a key regulator of connective tissue cells. *Kidney Int.* **51**, 1361-1369 (1997)
- Lindahl, P., Johansson, B., Levéen, P., and Betsholtz, C. Pericyte loss and microaneurysm development in PDGF-B deficient mice. *Science*, **277**, 242-245 (1997)
- Lindahl, P., Karlsson, L., Hellström, M., Gebre-Medhin, S., Willetts, K., Heath, J.K. and Betsholtz, C. Alveogenesis failure in PDGF-A deficient mice is coupled to lack of distal spreading of alveolar smooth muscle cell progenitors during lung development. *Development*, **124**, 3943-3953 (1997)
- Pekny, M., Stanness, K.A., Eliasson, C., Betsholtz, C. and Janigro, D. Impaired induction of blood-brain barrier properties in aortic endothelial cells by astrocytes from GFAP-deficient mice. *Glia*, in press (1997)

Gebre-Medhin, S., Mulder, H., Pekny, M., Zhang, Y.-Z., Törnelli, J., Westermark, P., Sundler, F., Ahren, B., and Betsholtz, C. Enhanced insulin secretion and action, obesity, and altered nociception in islet amyloid polypeptide/amylin null mice. *EMBO J.*, submitted (1998)

Calver, A.R., Hall, A.C., Walsh, F.S., Heath, J.K., Betsholtz, C. and Richardson, W.D. Control of glial progenitor cell number by competition for limiting mitogens. *Neuron*, submitted, (1998)

Falk, P., Betsholtz, C., Levéen, P., Lindahl, P., Franklin, G., Åhrlund-Richter, L., Pollard, J., Soriano, P., and Ohlsson, R. PDGF-B is required for the formation of an efficient nutrient-waste exchange system in the mouse fetal placenta. Submitted to *Development* (1997).

Kaminski, W., Lindahl, P., Crosby, J.R., Swolin, B., Bowen-Pope, D.F., Betsholtz, C., Martin, P., Ross, R., Raines, E. Chimeric mice deficient in PDGF-B or PDGF- β receptor in their hematologic cells are viable and display normal hematopoiesis. *J. Clin. Invest.* submitted (1998)

Pekna, M., Hietala, M.A., Landin, A., Betsholtz, C. and Pekny, M. Mice deficient for the complement factor B develop and reproduce normally. *Eur J Immunol.* Submitted (1998)

Pekna, M., Hietala, M.A., Rosklint, T., Betsholtz, C. and Pekny, M. Targeted disruption of the murine gene coding for the third complement component (C3). *Scand. J. Immunol.* in press (1997)

Lindal, P., Betsholtz, C. Not all myofibroblasts are alike or Revisiting the role of PDGF-AA and PDGF-BB using PDGF targeted mice. *Curr. Op. Nephrol. Hypertension.* In press 1998